

ET



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 413 528 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90308828.4

(51) Int. Cl.⁵: A61K 7/48, A61K 31/19

(22) Date of filing: 10.08.90

(30) Priority: 15.08.89 US 393749

(43) Date of publication of application:
20.02.91 Bulletin 91/08

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: Yu, Ruey J., Dr.
4 Lindenwold Avenue
Ambler Pennsylvania 19002(US)

Applicant: Van Scott, Eugene J., Dr.
3 Hidden Land
Abington Pennsylvania 19001(US)

(72) Inventor: Yu, Ruey J., Dr.
4 Lindenwold Avenue
Ambler Pennsylvania 19002(US)
Inventor: Van Scott, Eugene J., Dr.
3 Hidden Land
Abington Pennsylvania 19001(US)

(74) Representative: Bankes, Stephen Charles
Digby et al
BARON & WARREN 18 South End Kensington
London W8 5BU(GB)

(54) Amphoteric compositions and polymeric forms of alpha hydroxyacids, and their therapeutic use.

(57) Preventive as well as therapeutic treatment to alleviate cosmetic conditions and symptoms of dermatologic disorders with amphoteric compositions containing alpha hydroxyacids, alpha ketoacids, related compounds or polymeric forms of hydroxyacids is disclosed. The cosmetic conditions and the dermatologic disorders in which the amphoteric compositions and the polymeric compounds may be useful include dry skin, dandruff, acne, keratoses, psoriasis, eczema, pruritus, age spots, lentigines, melasmas, wrinkles, warts, blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, skin changes associated with aging, and skin requiring cleansers.

EP 0 413 528 A1

AMPHOTERIC COMPOSITIONS AND POLYMERIC FORMS OF ALPHA HYDROXYACIDS, AND THEIR THERAPEUTIC USE

This invention relates generally to therapeutic treatment as well as preventive measures for cosmetic conditions and dermatologic disorders by topical administration of amphoteric compositions or polymeric forms of alpha hydroxyacids, alpha ketoacids and related compounds. We initially discovered that alpha hydroxy or keto acids and their derivatives were effective in the topical treatment of disease conditions such as dry skin, ichthyosis, eczema, palmar and plantar hyperkeratoses, dandruff, acne and warts.

We have now discovered that amphoteric compositions and polymeric forms of alpha hydroxyacids, alpha ketoacids and related compounds on topical administration are therapeutically effective for various cosmetic conditions and dermatologic disorders.

In order prior U.S. Patent No. 3,879,537 entitled "Treatment of Ichthyosiform Dermatoses" we described and claimed the use of certain alpha hydroxyacids, alpha ketoacids and related compounds for topical treatment of fish-scale like ichthyotic conditions in humans. In our U.S. Patent No. 3,920,835 entitled "Treatment of Disturbed Keratinization" we described and claimed the use of these alpha hydroxyacids, alpha ketoacids and their derivatives for topical treatment of dandruff, acne, and palmar and plantar hyperkeratosis.

In our prior U.S. Patent No. 4,105,783 entitled "Treatment of Dry Skin" we described and claimed the use of alpha hydroxyacids, alpha ketoacids and their derivatives for topical treatment of dry skin. In our recent U.S. Patent No. 4,246,261 entitled "Additives Enhancing Topical Corticosteroid Action" we described and claimed that alpha hydroxyacids, alpha ketoacids and their derivatives, could greatly enhance the therapeutic efficacy of corticosteroids in topical treatment of psoriasis, eczema, seborrheic dermatitis and other inflammatory skin conditions.

In our more recent U.S. Patent No. 4,363,815 entitled "Alpha Hydroxyacids, Alpha Ketoacids and Their Use in Treating Skin Conditions" we described and claimed that alpha hydroxyacids and alpha ketoacids related to or originating from amino acids, whether or not found in proteins, were effective in topical treatment of skin disorders associated with disturbed keratinization or inflammation. These skin disorders include dry skin, ichthyosis, palmar and plantar hyperkeratosis, dandruff, Darier's disease, lichen simplex chronicus, keratoses, acne, psoriasis, eczema, pruritus, warts and herpes.

In our most recent patent application Serial No. 945,680 filed December 23, 1986 and entitled "Additives Enhancing Topical Actions of Therapeutic Agents" we described and claimed that incorporation of an alpha hydroxyacid or related compound can substantially enhance therapeutic actions of cosmetic and pharmaceutical agents.

There is no doubt that alpha hydroxyacids, alpha ketoacids and related compounds are therapeutically effective for topical treatment of various cosmetic conditions and dermatologic disorders including dry skin, acne, dandruff, keratoses, age spots, wrinkles and disturbed keratinization. However, the compositions containing these acids may irritate human skin on repeated topical applications due to lower pH of the formulations. The irritation may range from a sensation of tingling, itching and burning to clinical signs of redness and peeling. Causes for such irritation may arise from the following:

Upper layers of normal skin have a pH of 4.2 to 5.6, but the compositions containing most alpha hydroxyacids or alpha ketoacids have pH values of less than 3.0. For example, a topical formulation containing 7.6% (1 M) glycolic acid has a pH of 1.9, and a composition containing 9% (1 M) lactic acid has the same pH of 1.9. These compositions of lower pH on repeated topical applications can cause a drastic pH decrease in the stratum corneum of human skin, and provoke disturbances in intercorneocyte bondings resulting in adverse skin reactions, especially to some individuals with sensitive skin.

Moreover, with today's state of the art it is still very difficult to formulate a lotion, cream or ointment emulsion which contains a free acid form of the alpha hydroxyacid, and which is physically stable as a commercial product for cosmetic or pharmaceutical use.

When a formulation containing an alpha hydroxyacid or alpha ketoacid is reacted equimolarly or equinormally with a metallic alkali such as sodium hydroxide or potassium hydroxide the composition becomes therapeutically ineffective. The reasons for such loss of therapeutic effects are believed to be as follows:

The intact skin of humans is a very effective barrier to many natural and synthetic substances. Cosmetic and pharmaceutical agents may be pharmacologically effective by oral or other systematic administration, but many of them are much less or totally ineffective on topical application to the skin. Topical effectiveness of a pharmaceutical agent depends on two major factors; (a) bioavailability of the active ingredient in the topical preparation and (b) percutaneous absorption, penetration and distribution of

the active ingredient to the target site in the skin. For example, a topical preparation containing 5% salicylic acid is therapeutically effective as a keratolytic, but that containing 5% sodium salicylate is not an effective product. The reason for such difference is that salicylic acid is in bioavailable form and can penetrate the stratum corneum, but sodium salicylate is not in bioavailable form and cannot penetrate the stratum corneum of the skin.

In the case of alpha hydroxyacids, a topical preparation containing 5% glycolic acid is therapeutically effective for dry skin, but that containing 5% sodium glycollate is not effective. The same is true in case of 5% lactic acid versus 5% sodium lactate. The reason for such difference is that both glycolic acid and lactic acid are in bioavailable forms and can readily penetrate the stratum corneum, but sodium glycollate and sodium lactate are not in bioavailable forms and cannot penetrate the stratum corneum of the skin.

When a formulation containing an alpha hydroxyacid or alpha ketoacid is reacted equimolarly or equinormally with ammonium hydroxide or an organic base of smaller molecule the composition still shows some therapeutic effects for certain cosmetic conditions such as dry skin, but the composition has lost most of its potency for other dermatologic disorders such as wrinkles, keratoses, age spots and skin changes associated with aging.

It has now been discovered that amphoteric compositions containing alpha hydroxyacids, alpha ketoacids or related compounds, and also the compositions containing dimeric or polymeric forms of hydroxyacids overcome the aforementioned shortcomings and retain the therapeutic efficacies for cosmetic conditions and dermatologic disorders. The amphoteric composition contains in combination an amphoteric or pseudoamphoteric compound and at least one of the alpha hydroxyacids, alpha ketoacids or related compounds. Such amphoteric system has a suitable pH, and can release the active form of an alpha hydroxyacid or alpha ketoacid into the skin. The dimeric and polymeric forms of alpha, beta or other hydroxyacids in non-aqueous compositions have a more desired pH than that of the monomeric form of the hydroxyacids. The non-aqueous compositions can be formulated and induced to release the active form of hydroxyacids after the compositions have been topically applied to the skin. The cosmetic conditions and dermatologic disorders in humans and animals, in which the amphoteric compositions containing the dimeric or polymeric forms of hydroxyacids may be useful, include dry skin, dandruff, acne, keratoses, psoriasis, eczema, pruritus, age spots, lentigines, melasmas, wrinkles, warts, blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, skin changes associated with aging and as skin cleansers.

I. Amphoteric and Pseudoamphoteric Compositions

Amphoteric substances by definition should behave either as an acid or a base, and can be an organic or an inorganic compound. The molecule of an organic amphoteric compound should consist of at least one basic and one acidic group. The basic groups include, for example, amino, imino and guanido groups. The acidic groups include, for example, carboxylic, phosphoric and sulfonic groups. Some examples of organic amphoteric compounds are amino acids, peptides, polypeptides, proteins, creatine, aminoaldonic acids, aminouronic acids, lauryl aminopropylglycine, aminoaldaric acids, neuraminic acid, desulfated heparin, deacetylated hyaluronic acid, hyalobiuronic acid, chondrosine and deacetylated chondroitin.

Inorganic amphoteric compounds are certain metallic oxides such as aluminum oxide and zinc oxide.

Pseudoamphoteric compounds are either structurally related to true amphoteric compounds or capable of inducing the same function when they are incorporated into the compositions containing alpha hydroxyacids or ketoacids. Some examples of pseudoamphoteric compounds are creatinine, stearamidoethyl diethylamine, stearamidoethyl diethanolamine, stearamidopropyl dimethylamine, quaternary ammonium hydroxide and quaternium hydroxide.

The amphoteric composition of the instant invention contains in combination an alpha hydroxyacid or alpha ketoacid and an amphoteric or pseudoamphoteric compound. There are two advantages of utilizing an amphoteric or the like compound in the therapeutic composition containing an alpha hydroxy or ketoacid. These are (a) the overall pH of the composition is raised, so that the composition becomes less or non-irritating to the skin and (b) some alpha hydroxy or ketoacid molecules react with the amphoteric compound to form a quadruple ionic complex which acts as buffering system to control the release of alpha hydroxy or ketoacid into the skin, therefore, eliminating the skin irritation and still retaining the therapeutic efficacies.

The following are some examples. 2-Hydroxyethanoic acid (glycolic acid) 1 M aqueous solution has pH 1.9. The pHs of compositions change to 3.0 and 3.2 when arginine 0.5 M and creatinine 0.5 M respectively are incorporated into the formulations. 2-Hydroxypropanoic acid (lactic acid) 1 M aqueous solution has pH 1.9. The pHs of compositions change to 3.1 and 6.9 when arginine 0.5 M and 1.0 M respectively are

incorporated into the formulations. 2-Methyl 2-hydroxypropanoic acid (methylactic acid) 1 M aqueous solution has pH 1.9. The pHs of compositions change to 3.3, 3.4 and 3.2 when 0.5 M each of arginine, creatinine and 4-aminobutanoic acid respectively are incorporated into the formulations. 2-Hydroxybutane-1,4-dioic acid (malic acid) 1 M aqueous solution has pH 1.8, but the pH of the composition changes to 3.0 when creatinine 0.5 M is incorporated into the formulation.

Ideally, an amphoteric compound should contain both anionic and cationic groups or functional groups capable of behaving both as an acid and a base. Although inorganic amphoteric compounds such as aluminum oxide, aluminum hydroxide and zinc oxide may be utilized, organic amphoteric compounds have been found to be more efficient in formulating therapeutic compositions of the instant invention.

Organic amphoteric and pseudoamphoteric compounds may be classified into three groups, namely (a) amino acid type, (b) imidazoline and lecithin amphoteric and (c) pseudoamphoterics and miscellaneous amphoteric.

(a) Amino acid type amphoteric. Amphoteric compounds of amino acid type include all the amino acids, dipeptides, polypeptides, proteins and the like which contain at least one of the basic groups such as amino, imino, guanido, imidazolino and imidazolyl, and one of the acidic groups such as carboxylic, sulfonic, sulfinic and sulfate.

Glycine is a simple amphoteric compound which contains only one amino group and one carboxylic group. Lysine contains two amino groups and one carboxylic group. Aspartic acid contains one amino group and two carboxylic groups. Arginine contains one amino group, one guanido group and one carboxylic group. Histidine contains one amino group, one imidazolyl group and one carboxylic group. Taurine contains one amino group and one sulfonic group. Cysteine sulfinic acid contains one amino group, one carboxylic group and one sulfinic group. The amino group of an amphoteric compound may also be substituted, such as in betaine which is a glycine N,N,N-trimethyl inner salt.

Glycylglycine is a simple dipeptide which contains one free amino group and one free carboxylic group. Glycylhistidine is also a dipeptide which contains one free amino group, one imidazolyl group and one free carboxylic group.

The representative amphoteric compounds of amino acid type may be listed as follows: Glycine, alanine, valine, leucine, isoleucine, serine, threonine, cysteine, cystine, methionine, aspartic acid, asparagine, glutamic acid, glutamine, arginine, lysine, 5-hydroxylysine, histidine, phenylalanine, tyrosine, tryptophan, 3-hydroxyproline, 4-hydroxyproline and proline.

The related amino acids include homocysteine, homocystine, homoserine, ornithine, citrulline, creatine, 3-aminopropanoic acid, theanine, 2-aminobutanoic acid, 4-aminobutanoic acid, 2-amino-2-methylpropanoic acid, 2-methyl-3-aminopropanoic acid, 2,6-diaminopimelic acid, 2-amino-3-phenylbutanoic acid, phenylglycine, canavanine, cananine, 4-hydroxyarginine, 4-hydroxyornithine, homoarginine, 4-hydroxyhomoarginine, β -lysine, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid, 2-methylserine, 3-phenylserine and betaine.

Sulfur-containing amino acids include taurine, cysteinesulfinic acid, methionine sulfoxide and methionine sulfone.

The halogen-containing amino acids include 3,5-diiodotyrosine, thyroxine and monoiodotyrosine. The imino type acids include pipercolic acid, 4-aminopipercolic acid and 4-methylproline.

The dipeptides include for example, glycylglycine, carnosine, anserine, ophidine, homocarnosine, β -alanyllysine, β -alanylarginine. The tripeptides include for example, glutathione, ophthalmic acid and norophthalmic acid. Short-chain polypeptides of animal, plant and bacterial origin containing up to 100 amino acid residues include bradykinin and glucagon. The preferred proteins include for example protamines, histones and other lysine and arginine rich proteins.

(b) Imidazoline and lecithin amphoteric. The amphoteric compounds of imidazoline derived type are commercially synthesized from 2-substituted-2-imidazolines obtained by reacting a fatty acid with an aminoethylethanolamine. These amphoteric include cocoamphoglycine, cocoamphopropionate, and cocoamphopropylsulfonate. The amphoteric compounds of lecithin and related type include for example, phosphatidyl ethanolamine, phosphatidyl serine and sphingomyelin.

(c) Pseudoamphoterics and miscellaneous amphoteric. Many pseudoamphoteric compounds are chemically related or derived from true amphoteric. For example, creatinine is derived from creatine. Other pseudoamphoteric compounds may include fatty amide amines such as stearamidoethyl diethylamine, stearamidoethyl diethanolamine and stearamidopropyl dimethylamine. Other pseudoamphoteric related compounds include quaternary ammonium hydroxide and quaternium hydroxide.

In accordance with the present invention, the alpha hydroxyacid, the alpha ketoacids and the related compounds which are incorporated into amphoteric or pseudoamphoteric compositions for cosmetic conditions and dermatologic disorders may be classified into three groups.

The first group is organic carboxylic acids in which one hydroxyl group is attached to the alpha carbon

of the acids. The generic structure of such alpha hydroxyacids may be represented as follows:

(Ra) (Rb) C (OH) COOH

where Ra and Rb are H, F, Cl, Br, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and in addition Ra and Rb may carry OH, CHO, COOH and alkoxy group having 1 to 9 carbon atoms. The alpha hydroxyacids may be present as a free acid or lactone form, or in a salt form with an organic base or an inorganic alkali. The alpha hydroxyacids may exist as stereoisomers as D, L, and DL forms when Ra and Rb are not identical.

Typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, lauryl, stearyl, benzyl and phenyl, etc. The alpha hydroxyacids of the first group may be divided into (1) alkyl alpha hydroxyacids, (2) aralkyl and aryl alpha hydroxyacids, (3) polyhydroxy alpha hydroxyacids, and (4) polycarboxylic alpha hydroxyacids. The following are representative alpha hydroxyacids in each subgroup.

(1) Alkyl Alpha Hydroxyacids

1. 2-Hydroxyethanoic acid (Glycolic acid, hydroxyacetic acid)
(H) (H) C (OH) COOH
2. 2-Hydroxypropanoic acid (Lactic acid)
(CH₃) (H) C (OH) COOH
3. 2-Methyl 2-hydroxypropanoic acid (Methylactic acid)
(CH₃) (CH₃) C (OH) COOH
4. 2-Hydroxybutanoic acid
(C₂H₅) (H) C (OH) COOH
5. 2-Hydroxypentanoic acid
(C₃H₇) (H) C (OH) COOH
6. 2-Hydroxyhexanoic acid
(C₄H₉) (H) C (OH) COOH
7. 2-Hydroxyheptanoic acid
(C₅H₁₁) (H) C (OH) COOH
8. 2-Hydroxyoctanoic acid
(C₆H₁₃) (H) C (OH) COOH
9. 2-Hydroxynonanoic acid
(C₇H₁₅) (H) C (OH) COOH
10. 2-Hydroxydecanoic acid
(C₈H₁₇) (H) C (OH) COOH
11. 2-Hydroxyundecanoic acid
(C₉H₁₉) (H) C (OH) COOH
12. 2-Hydroxydodecanoic acid (Alpha hydroxylauric acid)
(C₁₀H₂₁) (H) C (OH) COOH
13. 2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid)
(C₁₂H₂₅) (H) C (OH) COOH
14. 2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid)
(C₁₄H₂₉) (H) C (OH) COOH
15. 2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid)
(C₁₆H₃₃) (H) C (OH) COOH
16. 2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid)
(C₁₈H₃₇) (H) C (OH) COOH

(2) Aralkyl And Aryl Alpha Hydroxyacids

1. 2-Phenyl 2-hydroxyethanoic acid (Mandelic acid)
(C₆H₅) (H) C (OH) COOH
2. 2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid)
(C₆H₅) (C₆H₅) C (OH) COOH
3. 3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid)
(C₆H₅CH₂) (H) C (OH) COOH
4. 2-Phenyl 2-methyl 2-hydroxyethanoic acid

- (Atrolactic acid)
 $(C_6H_5)(CH_3)C(OH)COOH$
 5. 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid
 (4-Hydroxymandelic acid)
 5 $(HO-C_6H_4)(H)C(OH)COOH$
 6. 2-(4'-Chlorophenyl) 2-hydroxyethanoic acid
 (4-Chloromandelic acid)
 $(Cl-C_6H_4)(H)C(OH)COOH$
 7. 2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic acid
 10 (3-Hydroxy-4-methoxymandelic acid)
 $(HO,CH_3O-C_6H_3)(H)C(OH)COOH$
 8. 2-(4'-Hydroxy-3'-methoxyphenyl) 2-hydroxyethanoic acid
 (4-Hydroxy-3-methoxymandelic acid)
 $(HO,CH_3O-C_6H_3)(H)C(OH)COOH$
 15 9. 3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid
 [3-(2'-Hydroxyphenyl) lactic acid]
 $HO-C_6H_4-CH_2(H)C(OH)COOH$
 10. 3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid
 [3-(4'-Hydroxyphenyl) lactic acid]
 20 $HO-C_6H_4-CH_2(H)C(OH)COOH$
 11. 2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid
 (3,4-Dihydroxymandelic acid)
 $HO,HO-C_6H_3(H)C(OH)COOH$
- 25 (3) Polyhydroxy Alpha Hydroxyacids
1. 2,3-Dihydroxypropanoic acid (Glyceric acid)
 $(HOCH_2)(H)C(OH)COOH$
 30 2. 2,3,4-Trihydroxybutanoic acid (Isomers; erythronic acid, threonic acid)
 $HOCH_2(HO)CH_2(H)C(OH)COOH$
 3. 2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic acid, arabinoic acid, xylonic acid, lyxonic acid)
 $HOCH_2(HO)CH_2(HO)CH_2(H)C(OH)COOH$
 4. 2,3,4,5,6-Pentahydroxyhexanoic acid (Isomers; allonic acid, altronic acid, gluconic acid, mannoic acid,
 gulonic acid, idonic acid, galactonic acid, talonic acid)
 35 $HOCH_2(HO)CH_2(HO)CH_2(HO)CH_2(H)C(OH)COOH$
 5. 2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid, galactoheptonic acid etc.)
 $HOCH_2(HO)CH_2(HO)CH_2(HO)CH_2(HO)CH_2(H)C(OH)COOH$
- (4) Polycarboxylic Alpha Hydroxyacids
- 40 1. 2-Hydroxypropane-1,3-dioic acid (Tartronic acid)
 $HOOC(H)C(OH)COOH$
 2. 2-Hydroxybutane-1,4-dioic acid (Malic acid)
 $HOOCCH_2(H)C(OH)COOH$
 45 3. 2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid)
 $HOOC(HO)CH(H)C(OH)COOH$
 4. 2-Hydroxy-2-carboxypentane-1,5-dioic acid (Citric acid)
 $HOOCCH_2C(OH)(COOH)CH_2COOH$
 5. 2, 3, 4, 5-Tetrahydroxyhexane- 1, 6-dioic acid (Isomers; saccharic acid, mucic acid etc.)
 50 $HOOC(CHOH)_4COOH$
- (5) Lactone Forms

The typical lactone forms are gluconolactone, galactonolactone, glucuronolactone, galacturonolactone,
 55 gulonolactone, ribonolactone, saccharic acid lactone, pantoyllactone, glucoheptonolactone, mannolactone,
 and galactoheptonolactone.

The second group of compounds which may be incorporated into amphoteric or pseudoamphoteric
 compositions for cosmetic conditions and dermatologic disorders, is organic carboxylic acids in which the

alpha carbon of the acids is in keto form. The generic structure of such alpha ketoacids may be represented as follows:

(Ra) CO COO(Rb)

wherein Ra and Rb are H, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and in addition Ra may carry F, Cl, Br, I, OH, CHO, COOH and alkoxy group having 1 to 9 carbon atoms. The alpha ketoacids may be present as a free acid or an ester form, or in a salt form with an organic base or an inorganic alkali. The typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, lauryl, stearyl, benzyl and phenyl, etc.

In contrast to alpha hydroxyacids the ester form of alpha ketoacids has been found to be therapeutically effective for cosmetic and dermatologic conditions and disorders. For example, while ethyl lactate has a minimal effect, ethyl pyruvate is therapeutically very effective. Although the real mechanism for such difference is not known, we have speculated that the ester form of an alpha ketoacid is chemically and/or biochemically very reactive, and a free acid form of the alpha ketoacid is released in the skin after the topical application.

The representative alpha ketoacids and their esters which may be useful in amphoteric or pseudoamphoteric compositions for cosmetic conditions and dermatologic disorders are listed below:

1. 2-Ketoethanoic acid (Glyoxylic acid)
(H) CO COOH
2. Methyl 2-ketoethanoate
(H) CO COOCH₃
3. 2-Ketopropanoic acid (Pyruvic acid)
CH₃ CO COOH
4. Methyl 2-ketopropanoate (Methyl pyruvate)
CH₃ CO COOCH₃
5. Ethyl 2-ketopropanoate (Ethyl pyruvate)
CH₃ CO COOC₂H₅
6. Propyl 2-ketopropanoate (Propyl pyruvate)
CH₃ CO COOC₃H₇
7. 2-Phenyl-2-ketoethanoic acid (Benzoylformic acid)
C₆H₅ CO COOH
8. Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate)
C₆H₅ CO COOCH₃
9. Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate)
C₆H₅ CO COOC₂H₅
10. 3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid)
C₆H₅CH₂ CO COOH
11. Methyl 3-phenyl-2-ketopropanoate (Methyl phenylpyruvate)
C₆H₅CH₂ CO COOCH₃
12. Ethyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate)
C₆H₅CH₂ CO COOC₂H₅
13. 2-Ketobutanoic acid
C₂H₅ CO COOH
14. 2-Ketopentanoic acid
C₃H₇ CO COOH
15. 2-Ketohexanoic acid
C₄H₉ CO COOH
16. 2-Ketoheptanoic acid
C₅H₁₁ CO COOH
17. 2-Ketooctanoic acid
C₆H₁₃ CO COOH
18. 2-Ketododecanoic acid
C₁₀H₂₁ CO COOH
19. Methyl 2-ketooctanoate
C₆H₁₃ CO COOCH₃

The third group of compounds which may be incorporated into amphoteric or pseudoamphoteric compositions for cosmetic and dermatologic conditions and disorders, is chemically related to alpha hydroxyacids or alpha ketoacids, and can be represented by their names instead of the above two generic

structures. The third group of compounds include ascorbic acid, quinic acid, isocitric acid, tropic acid, trethocanic acid, 3-chlorolactic acid, cerebronic acid, citramalic acid, agaricic acid, 2-hydroxynervonic acid, aleuritic acid and pantoic acid.

5

II. Dimeric and Polymeric Forms of Hydroxyacids

When two or more molecules of hydroxycarboxylic acids either identical or non-identical compounds are reacted chemically to each other, dimeric or polymeric compounds will be formed. Such dimeric and polymeric compounds may be classified into three groups, namely (a) acyclic ester, (b) cyclic ester and (c) miscellaneous dimer and polymer.

(a) Acyclic ester. The acyclic ester of a hydroxycarboxylic acid may be a dimer or a polymer. The dimer is formed from two molecules of a hydroxycarboxylic acid by reacting the carboxyl group of one molecule with the hydroxy group of a second molecule. For example, glycolyl glycollate is formed from two molecules of glycolic acid by eliminating one mole of water molecule. Likewise, lactyl lactate is formed from two molecules of lactic acid. When two molecules of different hydroxycarboxylic acids are intermolecularly reacted, a different dimer is formed. For example, glycolyl lactate is formed by reacting the carboxyl group of lactic acid with the hydroxy group of glycolic acid. The polymer is formed in a similar manner but from more than two molecules of a hydroxycarboxylic acid. For example, glycoly glycoly glycollate is formed from three molecules of glycolic acid. Copolymer is formed from two or more than two different kinds of hydroxycarboxylic acids. For example, glycolyl lactyl glycollate is formed from two molecules of glycolic acid and one molecule of lactic acid.

The acyclic ester of dimeric and polymeric hydroxycarboxylic acids may be shown by the following chemical structure:



wherein Ra, Rb = H, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and n = 1 or any numerical number, with a preferred number of up to 200. Ra and Rb in monomer unit 2, 3, 4 and so on may be the same or the different groups from that in monomer unit 1. For example, Ra, Rb = H in monomer unit 1, and Ra = CH₃, Rb = H in monomer unit 2 when n = 2 is a dimer called lactyl glycollate, because the first monomer is glycollate unit and the second monomer is lactic acid unit. The hydrogen atom in Ra and Rb may be substituted by a halogen atom or a radical such as a lower alkyl, aralkyl, aryl or alkoxy of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 9 carbon atoms. The dimer and polymer of a hydroxycarboxylic acid may be present as a free acid, ester or salt form with organic base or inorganic alkali.

The typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, benzyl and phenyl. Representative acyclic esters of hydroxycarboxylic acids which may be useful for cosmetic conditions and dermatologic disorders are listed below:

1. Glycolyl glycollate (Glycolic acid glycollate)
Ra, Rb = H in units 1 & 2, n = 2
2. Lactyl lactate (Lactic acid lactate)
Ra = CH₃, Rb = H in units 1 & 2, n = 2
3. Mandelyl mandellate
Ra = C₆H₅, Rb = H in units 1 & 2, n = 2
4. Atrolactyl atrolactate
Ra = C₆H₅, Rb = CH₃ in units 1 & 2, n = 2
5. Phenyllactyl phenyllactate
Ra = C₆H₅CH₂, Rb = H, in units 1 & 2, n = 2
6. Benzilyl benzillate
Ra, Rb = C₆H₅ in units 1 & 2, n = 2
7. Glycolyl lactate
Ra = CH₃ in unit 1, Ra = H in unit 2, Rb = H in units 1 & 2, n = 2
8. Lactyl glycollate
Ra = H in unit 1, Ra = CH₃ in unit 2, Rb = H in units 1 & 2, n = 2
9. Glycolyl glycolyl glycollate
Ra, Rb = H in units 1, 2 & 3, n = 3
10. Lactyl lactyl lactate
Ra = CH₃, Rb = H in units 1, 2 & 3, n = 3

11. Lactyl glycolyl lactate
 $Ra=CH_3$ in units 1 & 3, $Ra=H$ in unit 2, $Ra=H$ in units 1, 2 & 3, $n=3$
12. Glycolyl glycolyl glycolyl glycollate
 $Ra,Rb=H$ in units 1, 2, 3 & 4, $n=4$
13. Lactyl lactyl lactyl lactate
 $Ra=CH_3$, $Rb=H$ in units 1, 2, 3 & 4, $n=4$
14. Glycolyl lactyl glycolyl lactyl glycollate
 $Ra=H$ in units 1, 3 & 5, $Ra=CH_3$ in units 2 & 4, $Rb=H$ in units 1, 2, 3, 4 & 5, $n=5$
15. Polyglycolic acid and polylactic acid
- (b) Cyclic ester. The cyclic ester of a hydroxycarboxylic acid may also be a dimer or polymer, the most common type however, is a dimer form. The cyclic dimer may be formed from an identical monomer or different monomers. For example, glycolide is formed from two molecules of glycolic acid by removing two molecules of water, and lactide is formed from two molecules of lactic acid in the same manner. The cyclic ester of dimeric and polymeric hydroxycarboxylic acids may be shown by the following chemical structure:
- $[-O-C(Ra)(Rb)-Co-]_n$
 wherein $Ra,Rb=H$, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and $n=1$ or any numerical number, however with a preferred number of 2. Ra and Rb in units 1, 2, 3 and so on may be the same or the different groups. For example, in glycolide Ra and Rb are H in both units 1 & 2, but in lactoglycolide Ra is H in unit 1, CH_3 in unit 2 and Rb is H in both units 1 & 2. The hydrogen atom in Ra and Rb may be substituted by a halogen atom or a radical such as a lower alkyl, aralkyl, aryl or alkoxy of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 9 carbon atoms.
- The typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, benzyl and phenyl. Representative cyclic esters of hydroxycarboxylic acids which may be useful for cosmetic conditions and dermatologic disorders are listed below:
1. Glycolide
 $Ra,Rb=H$, $n=2$
 2. Lactide
 $Ra=CH_3$, $Rb=H$ in units 1 & 2, $n=2$
 3. Mandelide
 $Ra=C_6H_5$, $Rb=H$ in units 1 & 2, $n=2$
 4. Atrolactide
 $Ra=C_6H_5$, $Rb=CH_3$ in units 1 & 2, $n=2$
 5. Phenyllactide
 $Ra=C_6H_5$, $Rb=CH_2$, $Rb=H$ in units 1 & 2, $n=2$
 6. Benzilide
 $Ra,Rb=C_6H_5$ in units 1 & 2, $n=2$
 7. Methyllactide
 $Ra,Rb=CH_3$ in units 1 & 2, $n=2$
 8. Lactoglycolide
 $Ra=H$ in unit 1, $Ra=CH_3$ in unit 2
 $Rb=H$ in units 1 & 2, $n=2$
 9. Glycolactide
 $Ra=CH_3$ in unit 1, $Ra=H$ in unit 2
 $Rb=H$ in units 1 & 2, $n=2$
- (c) Miscellaneous dimer and polymer. This group includes all the dimeric and polymeric forms of hydroxycarboxylic acids, which can not be represented by any one of the above two generic structures, such as those formed from tropic acid, trethocanic acid and aleuritic acid. When a hydroxycarboxylic acid has more than one hydroxy or carboxy group in the molecule a complex polymer may be formed. Such complex polymer may consist of acyclic as well as cyclic structures.
- The following hydroxycarboxylic acids have more than one hydroxy groups: glyceric acid, gluconic acid and gluconolactone, galactonic acid and galactonolactone, glucuronic acid and glucuronolactone, ribonic acid and ribonolactone, galacturonic acid and galacturonolactone, ascorbic acid, gulonic acid and gulonolactone, glucoheptonic acid and glucoheptonolactone. These polyhydroxycarboxylic acids can form complex polymers with themselves or with other simple monohydroxymonocarboxylic acids.
- The following hydroxycarboxylic acids have more than one carboxyl groups: malic acid, citric acid, citramalic acid, tartronic acid, agaricic acid and isocitric acid. These monohydroxypolycarboxylic acids can also form complex polymers with themselves or with other simple hydroxycarboxylic acids.
- The following hydroxycarboxylic acids have more than one hydroxy and more than one carboxyl groups: tartaric acid, mucic acid and saccharic acid. These polyhydroxypolycarboxylic acids can form even

more complex polymers with themselves or with other hydroxycarboxylic acids.

III. Combination Compositions

5

Any cosmetic and pharmaceutical agents may be incorporated into amphoteric or pseudoamphoteric compositions, or into compositions containing dimeric or polymeric forms of hydroxyacids with or without amphoteric or pseudoamphoteric systems to enhance therapeutic effects of those cosmetic and pharmaceutical agents to improve cosmetic conditions or to alleviate the symptoms of dermatologic disorder.

10 Cosmetic and pharmaceutical agents include those that improve or eradicate age spots, keratoses and wrinkles; analgesics; anesthetics; antiacne agents; antibacterials; antiyeast agents; antifungal agents; antiviral agents; antidandruff agents; antidermatitis agents; antipruritic agents; antiemetics; antimoion sickness agents; antiinflammatory agents; antihyperkeratolytic agents; antidryskin agents; antiperspirants; antipsoriatic agents; antiseborrheic agents; hair conditioners and hair treatment agents; antiaging and antiwrinkle agents; antiasthmatic agents and bronchodilators; sunscreen agents; antihistamine agents; skin lightening agents; depigmenting agents; vitamins; corticosteroids; tanning agents; hormones; retinoids; topical cardiovascular agents and other dermatologicals.

Some examples of cosmetic and pharmaceutical agents are clotrimazole, ketoconazole, miconazole, griseofulvin, hydroxyzine, diphenhydramine, pramoxine, lidocaine, procaine, mepivacaine, monobenzene, 20 erythromycin, tetracycline, clindamycin, meclocyline, hydroquinone, minocycline, naproxen, ibuprofen, theophylline, cromolyn, albuterol, retinoic acid, 13-cis retinoic acid, hydrocortisone, hydrocortisone 21-acetate, hydrocortisone 17-valerate, hydrocortisone 17-butyrate, betamethasone valerate, betamethasone dipropionate, triamcinolone acetonide, flucinonide, clobetasol propionate, benzoyl peroxide, crotamiton, propranolol, promethazine, vitamin A palmitate and vitamin E acetate.

25

IV. Specific Compositions For Skin Disorders

We have discovered that topical formulations or compositions containing specific alpha hydroxyacids or 30 alpha ketoacids, or related compounds are therapeutically very effective for certain skin disorders without utilizing any amphoteric or pseudoamphoteric systems. The alpha hydroxyacids and the related compounds include 2-hydroxyethanoic acid, 2-hydroxypropanoic acid, 2-methyl 2-hydroxypropanoic acid, 2-phenyl 2-hydroxyethanoic acid, 2,2-diphenyl 2-hydroxyethanoic acid, 2-phenyl 2-methyl 2-hydroxyethanoic acid and 2-phenyl 3-hydroxypropanoic acid. The alpha ketoacids and their esters include 2-ketopropanoic acid, 35 methyl 2-ketopropanoate and ethyl 2-ketopropanoate. The mentioned skin disorders include warts, keratoses, age spots, acne, nail infections, wrinkles and aging related skin changes.

In general, the concentration of the alpha hydroxyacid, the alpha ketoacid or the related compound used in the composition is a full strength to an intermediate strength, therefore the dispensing and the application require special handling and procedures.

40 If the alpha hydroxyacid, or the alpha ketoacid or the related compound at full strength (usually 95-100%) is a liquid form at room temperature such as 2-hydroxypropanoic acid, 2-ketopropanoic acid, methyl 2-ketopropanoate and ethyl 2-ketopropanoate, the liquid compound with or without a gelling agent is directly dispensed as 0.5 to 1 ml aliquots in small vials.

If the alpha hydroxyacid, or the alpha ketoacid or the related compound at full strength is a solid form at 45 room temperature such as 2-hydroxyethanoic acid, 2-methyl 2-hydroxypropanoic acid, 2-phenyl 2-hydroxyethanoic acid, 2,2-diphenyl 2-hydroxyethanoic acid and 2-phenyl 3-hydroxypropanoic acid, the solid compound is first dissolved in a minimal amount of vehicle or vehicle system such as water, or ethanol and propylene glycol with or without a gelling agent. For example, 2-hydroxyethanoic acid 70 g is dissolved in water 30 g, and the 70% strength solution thus obtained is dispensed as 0.5 to 1 ml aliquots in small vials. 50 If a gelling agent is used, 0.5 to 3% of for example, hydroxyethyl cellulose, methyl cellulose, hydroxypropyl cellulose or carbomer may be incorporated into the above solution.

To prepare an intermediate strength (usually 20-50%), the alpha hydroxyacid, alpha ketoacid or related compound either a liquid or solid form at room temperature is first dissolved in a vehicle or vehicle system such as water, acetone, ethanol, propylene glycol and butane 1,3-diol. For example, 2-hydroxyethanoic acid 55 or 2-ketopropanoic acid 30 g is dissolved in ethanol 56 g and propylene glycol 14 g, and the 30% strength solution thus obtained is dispensed as 7 to 14 ml aliquots in dropper bottles.

For topical treatment of warts, keratoses, age spots, acne, nail infections, wrinkles or aging related skin changes, patients are advised to apply a small drop of the medication with a toothpick or a fine-caliber,

commonly available artist's camel hair brush to affected lesions only and not surrounding skin. Prescribed applications have been 1 to 6 times daily for keratoses and ordinary warts of the hands, fingers, palms, and soles. For age spots, acne, nail infections, wrinkles and aging related skin changes topical applications have been 1 to 2 times daily.

5 Very often, frequency and duration of applications have been modified according to clinical responses and reactions of the lesions and the patient or responsible family member is instructed accordingly. For example, some clinical manifestations other than pain have been used as a signal to interrupt application. These manifestations include distinct blanching of the lesions or distinct peripheral erythema.

Alternatively, an office procedure may be adapted when a full strength of 2-ketopropanoic acid or 70%
10 2-hydroxyethanoic acid is used for topical treatment of age spots, keratoses, acne, warts or facial wrinkles.

We have found that the above mentioned alpha hydroxyacids, alpha ketoacids and related compounds are therapeutically effective for topical treatments of warts, keratoses, age spots, acne, nail infections, wrinkles and aging related skin changes.

15

Preparation of the Therapeutic Compositions

Amphoteric and pseudoamphoteric compositions of the instant invention may be formulated as solution,
20 gel, lotion, cream, ointment, shampoo, spray, stick, powder or other cosmetic and pharmaceutical preparations.

To prepare an amphoteric or pseudoamphoteric composition in solution form at least one of the aforementioned amphoteric or pseudoamphoteric compounds and in combination at least one of the hydroxyacids or the related compounds are dissolved in a solution which may consist of ethanol, water,
25 propylene glycol, acetone or other pharmaceutically acceptable vehicle. The concentration of the amphoteric or pseudoamphoteric compound may range from 0.01 to 10 M, the preferred concentration ranges from 0.1 to 3 M. The concentration of hydroxyacids or the related compounds may range from 0.02 to 12 M, the preferred concentration ranges from 0.2 to 5 M.

In the preparation of an amphoteric or pseudoamphoteric composition in lotion, cream or ointment form,
30 at least one of the amphoteric or pseudoamphoteric compounds and one of the hydroxyacids or the related compounds are initially dissolved in a solvent such as water, ethanol and/or propylene glycol. The solution thus prepared is then mixed in a conventional manner with commonly available cream or ointment base such as hydrophilic ointment or petrolatum. The concentrations of amphoteric or pseudoamphoteric compounds and hydroxyacids used in the compositions are the same as described above.

35 Amphoteric and pseudoamphoteric compositions of the instant invention may also be formulated in a gel form. A typical gel composition of the instant invention utilizes at least one of the amphoteric or pseudoamphoteric compounds and one of the hydroxyacids or the related compounds are dissolved in a mixture of ethanol, water and propylene glycol in a volume ratio of 40:40:20, respectively. A gelling agent such as methyl cellulose, ethyl cellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-
40 methylcellulose, carbomer or ammoniated glycyrrhizinate is then added to the mixture with agitation. The preferred concentration of the gelling agent may range from 0.1 to 4 percent by weight of the total composition.

Since dimeric and polymeric forms of hydroxyacids are less stable in the presence of water or the like vehicle, cosmetic and pharmaceutical compositions should be prepared as anhydrous formulations. Typical
45 vehicles suitable for such formulations include mineral oil, petrolatum, isopropyl myristate, isopropyl palmitate, diisopropyl adipate, octyl palmitate, acetone, squalene, squalane, silicone oils, vegetable oils and the like. Therapeutic compositions containing dimeric or polymeric forms of hydroxyacids do not require any incorporation of an amphoteric or pseudoamphoteric compound. The concentration of the dimeric or polymeric form of a hydroxyacid used in the composition may range from 0.1 to 100%, the
50 preferred concentration ranges from 1 to 40%. Therapeutic compositions may be formulated as anhydrous solution, lotion, ointment, spray, powder or the like.

To prepare a combination composition in a pharmaceutically acceptable vehicle, a cosmetic or pharmaceutical agent is incorporated into any one of the above composition by dissolving or mixing the agent into the formulation.

55 The following are illustrative examples of formulations and compositions according to this invention. Although the examples utilize only selected compounds and formulations, it should be understood that the following examples are illustrative and not limited. therefore, any of the aforementioned amphoteric or pseudoamphoteric compounds, hydroxyacids, dimeric or polymeric forms of hydroxyacids may be substi-

tuted according to the teachings of this invention in the following examples.

EXAMPLE 1

5

An amphoteric composition containing 1 M 2-hydroxyethanoic acid and 0.5 M L-arginine in solution form for dandruff or dry skin may be formulated as follows.

2-Hydroxyethanoic acid (glycolic acid) 7.6 g is dissolved in water 60 ml and propylene glycol 20 ml. L-Arginine 8.7 g is added to the solution with stirring until all the crystals are dissolved. Ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.0. An amphoteric composition formulated from 1 M 2-hydroxyethanoic acid and 1 M L-arginine has pH 6.3. The solution has pH 1.9 if no amphoteric compound is incorporated.

15

EXAMPLE 2

An amphoteric composition containing 1 M 2-hydroxyethanoic acid and 0.5 M L-lysine in a cream form for dry skin and other dermatologic and cosmetic conditions may be formulated as follows.

2-Hydroxyethanoic acid 7.6 g and L-lysine 7.3 g are dissolved in 30 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.3.

25

EXAMPLE 3

An amphoteric composition containing 1 M 2-hydroxyethanoic acid and 0.5 M 4-aminobutanoic acid in lotion form for cosmetic and dermatologic conditions may be formulated as follows.

2-Hydroxyethanoic acid 7.6 g and 4-aminobutanoic acid 5.2 g are dissolved in water 30 ml, and the solution is mixed with 50 g of an oil-in-water emulsion. The lotion thus obtained is made up to 100 ml in volume with more oil-in-water emulsion. The amphoteric composition thus formulated has pH 3.1.

35

EXAMPLE 4

A pseudoamphoteric composition containing 1 M 2-hydroxyethanoic acid and 0.5 M creatinine in solution form for cosmetic conditions and dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 7.6 g is dissolved in water 70 ml and propylene glycol 10 ml. Creatinine 5.7 g is added to the solution with stirring until all the crystals are dissolved. More water is added to make a total volume of the solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 3.2. The composition has pH 4.0 when 1 M instead of 0.5 M creatinine is incorporated into the formulation.

45

EXAMPLE 5

An amphoteric composition containing 1 M 2-hydroxyethanoic acid and 0.5 M L-histidine in a cream form for dermatologic and cosmetic conditions may be formulated as follows.

2-Hydroxyethanoic acid 7.6 g and L-histidine 7.8 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.2.

55

EXAMPLE 6

An amphoteric composition containing 0.5 M 2-hydroxyethanoic acid and 0.5 M dipeptide of β -Ala-L-His for cosmetic and dermatologic conditions may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and L-carnosine (β -alanyl-L-histidine) 11.3 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 4.5.

EXAMPLE 7

10

An amphoteric composition containing 0.5 M 2-hydroxyethanoic acid and 0.5 M cycloleucine for cosmetic and dermatologic conditions may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and 1-aminocyclopentane-1-carboxylic acid (cycloleucine) 6.5 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.2.

EXAMPLE 8

20

A pseudoamphoteric composition containing 0.5 M 2-hydroxyethanoic acid and 0.25 M 1,12-diaminododecane for cosmetic and dermatologic conditions may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and 1,12-diaminododecane 5 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

EXAMPLE 9

30

An amphoteric composition containing 0.5 M 2-hydroxyethanoic acid and 5% protamine for cosmetic and dermatologic conditions may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and protamine 5 g, isolated and purified from salmon sperm are dissolved in water 25 ml. The solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.2.

EXAMPLE 10

40

An amphoteric composition containing 1 M 2-hydroxypropanoic acid and 0.5 M L-arginine in solution form for dandruff or dry skin may be formulated as follows.

2-Hydroxypropanoic acid (DL-lactic acid) USP grade 9.0 g is dissolved in water 60 ml and propylene glycol 20 ml. L-Arginine 8.7 g is added to the solution with stirring until all the crystals are dissolved. Ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.1. An amphoteric composition formulated from 1 M 2-hydroxypropanoic acid and 1 M L-arginine has pH 6.9. The solution has pH 1.9 if no amphoteric compound is incorporated.

50

EXAMPLE 11

An amphoteric composition containing 1M 2-hydroxypropanoic acid and 0.5 M L-lysine in a cream form for dry skin and other dermatologic and cosmetic conditions may be formulated as follows.

2-Hydroxypropanoic acid 9.0 g and L-lysine 7.3 g are dissolved in 30 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The

55

amphoteric composition thus formulated has pH 3.6. An amphoteric composition formulated from 1 M 2-hydroxypropanoic acid and 1 M L-lysine has pH 8.4

5

EXAMPLE 12

An amphoteric composition containing 1 M 2-hydroxypropanoic acid and 0.5 M 4-aminobutanoic acid in lotion form for cosmetic and dermatologic conditions may be formulated as follows.

10 2-Hydroxypropanoic acid 9.0 g and 4-aminobutanoic acid 5.2 g are dissolved in water 30 ml, and the solution is mixed with 50 g of an oil-in-water emulsion. The lotion thus obtained is made up to 100 ml in volume with more oil-in-water emulsion. The amphoteric composition thus formulated has pH 3.0

15

EXAMPLE 13

A pseudoamphoteric composition containing 1 M 2-hydroxypropanoic acid and 0.5 M creatinine in solution form for cosmetic conditions and dermatologic disorders may be formulated as follows.

20 2-Hydroxypropanoic acid 9.0 g is dissolved in water 70 ml and propylene glycol 10 ml. Creatinine 5.7 g is added to the solution with stirring until all the crystals are dissolved. More water is added to make a total volume of the solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 3.3. The composition has pH 4.4 when 1 M instead of 0.5 M creatinine is incorporated into the formulation.

25

EXAMPLE 14

An amphoteric composition containing 1 M 2-hydroxypropanoic acid and 1 M L-histidine in a cream form for dermatologic and cosmetic conditions may be formulated as follows.

30 2-Hydroxypropanoic acid 9.0 g and L-histidine 15.5 g are dissolved in 35 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated as pH 4.9.

35

EXAMPLE 15

An amphoteric composition containing 1 M 2-hydroxypropanoic acid and 1 M dipeptide of Gly-Gly for cosmetic and dermatologic conditions may be formulated as follows.

40 2-Hydroxypropanoic acid 9.0 g and glycylglycine 13.2 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.0.

45

EXAMPLE 16

An amphoteric composition containing 1 M 2-methyl-2-hydroxypropanoic acid and 0.5 M L-arginine in solution form for dandruff or dry skin may be formulated as follows.

50 2-Methyl-2-hydroxypropanoic acid (methylactic acid) 10.4 g is dissolved in water 60 ml and propylene glycol 20 ml. L-Arginine 8.7 g is added to the solution with stirring until all the crystals are dissolved. Ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.3. An amphoteric composition formulated from 1 M 2-methyl-2-hydroxypropanoic acid
55 and 1 M L-arginine has pH 6.5. The solution has pH 1.9 if no amphoteric compound is incorporated.

EXAMPLE 17

An amphoteric composition containing 1 M 2-methyl-2-hydroxypropanoic acid and 0.5 M 4-aminobutanoic acid in a cream form for dry skin and other dermatologic and cosmetic conditions may be formulated as follows.

2-Methyl-2-hydroxypropanoic acid 10.4 g and 4-aminobutanoic acid 5.2 g are dissolved in 30 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.2.

EXAMPLE 18

An amphoteric composition containing 1 M 2-methyl-2-hydroxypropanoic acid and 1 M dipeptide of Gly-Gly in lotion form for cosmetic and dermatologic conditions may be formulated as follows.

2-Methyl-2-hydroxypropanoic acid 10.4 g and glycylglycine 13.2 g are dissolved in water 30 ml, and the solution is mixed with 50 g of an oil-in-water emulsion. The lotion thus obtained is made up to 100 ml in volume with more oil-in-water emulsion. The amphoteric composition thus formulated has pH 3.0.

EXAMPLE 19

A pseudoamphoteric composition containing 1 M 2-methyl-2-hydroxypropanoic acid and 0.5 M creatinine in solution form for cosmetic conditions and dermatologic disorders may be formulated as follows.

2-Methyl-2-hydroxypropanoic acid 10.4 g is dissolved in water 70 ml and propylene glycol 10 ml. Creatinine 5.7 g is added to the solution with stirring until all the crystals are dissolved. More water is added to make a total volume of the solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 3.4. The composition has pH 4.4 when 1 M instead of 0.5 M creatinine is incorporated into the formulation.

EXAMPLE 20

An amphoteric composition containing 0.5 M 2-phenyl-2-hydroxyethanoic acid and 0.5 M L-histidine in a cream form for dermatologic and cosmetic conditions may be formulated as follows.

2-Phenyl 2-hydroxyethanoic acid (mandelic acid) 7.6 g and L-histidine 7.8 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 5.0. The composition has pH 2.2 if no amphoteric compound is incorporated.

EXAMPLE 21

An amphoteric composition containing 0.5 M 2-phenyl-2-hydroxyethanoic acid and 0.5 M L-lysine for cosmetic and dermatologic conditions may be formulated as follows.

2-Phenyl 2-hydroxyethanoic acid 7.6 g and L-lysine 7.3 g are dissolved in 25 ml of water. The solution thus obtained is mixed with an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated for pH 4.6.

EXAMPLE 22

A pseudoamphoteric composition containing 0.5 M 2-phenyl 2-hydroxyethanoic acid and 0.5 M creatinine for cosmetic and dermatologic conditions may be formulated as follows.

2-Phenyl 2-hydroxyethanoic acid 7.6 g and creatinine 5.7 g are dissolved in 30 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 4.6.

5

EXAMPLE 23

An amphoteric composition containing 0.5 M 2-phenyl 2-hydroxyethanoic acid and 0.5 M L-citrulline for
10 cosmetic and dermatologic conditions may be formulated as follows.

2-Phenyl 2-hydroxyethanoic acid 7.6 g and L-citrulline 8.8 g are dissolved in water 30 ml, and the solution is mixed with 50 g of an oil-in-water emulsion. The lotion thus obtained is made up to 100 ml in volume with more oil-in-water emulsion. The amphoteric composition thus formulated has pH 3.0.

15

EXAMPLE 24

An amphoteric composition containing 1 M citric acid and 1 M L-arginine for cosmetic conditions and
20 dermatologic disorders may be formulated as follows.

Citric acid 19.2 g is dissolved in water 50 ml and propylene glycol 10 ml. L-Arginine 17.4 g is added to the solution with stirring until all the crystals are dissolved. More water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.0. The composition has pH 1.8 if no amphoteric compound is incorporated.

25

EXAMPLE 25

A pseudoamphoteric composition containing 1 M citric acid and 1 M creatinine for dermatologic and
30 cosmetic conditions may be formulated as follows.

Citric acid 19.2 g and creatinine 11.3 g are dissolved in 40 ml of water, and the solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.7.

35

EXAMPLE 26

An amphoteric composition containing 1 M malic acid and 1 M L-arginine for cosmetic and dermatologic
40 conditions may be formulated as follows.

2-Hydroxybutanedioic acid (DL-malic acid) 13.4 g and L-arginine 17.4 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.3. The
45 composition has pH 1.8 if no amphoteric compound is incorporated.

EXAMPLE 27

50

A pseudoamphoteric composition containing 1 M malic acid and 0.5 M creatinine for cosmetic and dermatologic conditions may be formulated as follows.

DL-Malic acid 13.4 g and creatinine 5.7 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of water is added to make a total volume of the
55 solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 3.0. The composition has pH 3.8 when 1 M instead of 0.5 M creatinine is incorporated into the formulation.

EXAMPLE 28

5 An amphoteric composition containing 1 M tartaric acid and 1 M L-arginine for cosmetic and dermatologic conditions may be formulated as follows.

2,3-Dihydroxybutanedioic acid (DL-tartaric acid) 15.9 g and L-arginine 17.4 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.0. The composition has pH 1.7 if no amphoteric compound is incorporated.

10

EXAMPLE 29

15 A pseudoamphoteric composition containing 1 M tartaric acid and 1 M creatinine for cosmetic and dermatologic conditions may be formulated as follows.

DL-Tartaric acid 15.0 g and creatinine 11.3 g are dissolved in 35 ml of water. The solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 3.4.

20

EXAMPLE 30

25 An amphoteric composition containing 1 M gluconolactone and 0.5 M L-arginine for cosmetic and dermatologic conditions may be formulated as follows.

Gluconolactone 17.8 g and L-arginine 8.7 g are dissolved in water 60 ml and propylene glycol 10 ml. After all the crystals have been dissolved sufficient water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.1. The composition has pH 5.9 when 1 M instead of 0.5 M L-arginine is incorporated into the formulation. If no amphoteric compound is incorporated the pH of the composition is 1.8.

30

EXAMPLE 31

35

An amphoteric composition containing 1 M gluconolactone and 0.5 M 4-aminobutanoic acid for cosmetic and dermatologic conditions may be formulated as follows.

Gluconolactone 17.8 g and 4-aminobutanoic acid 5.2 g are dissolved in water 60 ml and propylene glycol 10 ml. After all the crystals have been dissolved sufficient water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.2.

40

EXAMPLE 32

45

An amphoteric composition containing 1 M gluconolactone and 1 M dipeptide of Gly-Gly for cosmetic and dermatologic conditions may be formulated as follows.

Gluconolactone 17.8 g and glycylglycine 13.2 g are dissolved in water 50 ml and propylene glycol 5 ml. More water is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 3.1

50

EXAMPLE 33

55

A pseudoamphoteric composition containing 1 M gluconolactone and 0.5 M creatinine for cosmetic conditions and dermatologic disorders may be formulated as follows.

Gluconolactone 17.8 g and creatinine 5.7 g are dissolved in water 60 ml and propylene glycol 10 ml. More water is added to make a total volume of the solution to 100 ml. The pseudoamphoteric composition thus formulated has pH 3.2. The composition has pH 4.8 when 1 M instead of 0.5 M creatinine is incorporated into the formulation.

5

EXAMPLE 34

10 A pseudoamphoteric composition containing 1 M pyruvic acid and 1 M creatinine for dermatologic and cosmetic conditions may be formulated as follows.

2-Ketopropanoic acid (pyruvic acid) 8.8 g and creatinine 11.3 g are dissolved in water 25 ml. The solution thus obtained is mixed with sufficient amount of an oil-in-water emulsion to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 3.4.

15

EXAMPLE 35

20 An amphoteric composition containing 0.5 M benzoic acid and 0.5 M L-lysine for cosmetic and dermatologic conditions may be formulated as follows.

2,2-Diphenyl 2-hydroxyethanoic acid (benzoic acid) 11.4 g and L-lysine 7.3 g are dissolved in water 40 ml and propylene glycol 20 ml. After all the crystals have been dissolved sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 4.9. The composition has pH 2.7 if no amphoteric compound is incorporated.

25

EXAMPLE 36

30

An amphoteric composition containing 0.5 M benzoic acid and 0.5 M L-histidine for cosmetic and dermatologic conditions may be formulated as follows.

Benzoic acid 11.4 g and L-histidine 7.8 g are dissolved in water 40 ml and propylene glycol 20 ml. Ethyl cellulose 2 g is added with stirring, and sufficient amount of ethanol is added to make a total volume of the gel to 100 ml. The amphoteric gel composition thus formulated has pH 5.0.

35

EXAMPLE 37

40

A pseudoamphoteric composition containing 0.5 M benzoic acid and 0.5 M creatinine for cosmetic and dermatologic conditions may be formulated as follows.

Benzoic acid 11.4 g and creatinine 5.7 g are dissolved in water 40 ml and propylene glycol 20 ml. Sufficient amount of ethanol is added to make a total volume of the solution to 100 ml. The amphoteric composition thus formulated has pH 4.9.

45

EXAMPLE 38

50

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 0.05 % betamethasone dipropionate in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Betamethasone dipropionate 1 % in ethanol solution 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

55

EXAMPLE 39

5 A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 0.05% clobetasol propionate in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Clobetasol propionate 1 % in acetone solution 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

10

EXAMPLE 40

15 A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 0.1% triamcinolone acetonide in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Triamcinolone acetonide 2% solution of acetone:ethanol (50:50), 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

20

EXAMPLE 41

25 A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 0.2 % 5-fluorouracil in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g and creatinine 5.7 g are dissolved in 20 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. 5-Fluorouracil 2% solution of propylene glycol: water (95:5), 10 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

30

EXAMPLE 42

35 A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxypropanoic acid and 0.05 % betamethasone dipropionate in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxypropanoic acid 4.5 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Betamethasone dipropionate 1% in ethanol solution 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

40

EXAMPLE 43

45 A pseudoamphoteric composition containing in combination 0.5 M hydroxypropanoic acid and 0.05 % clobetasol propionate in a cream form for dermatologic disorders may be formulated as follows.

50 2-Hydroxypropanoic acid 4.5 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Clobetasol propionate 1% in acetone solution 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

55

EXAMPLE 44

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxypropanoic acid and 0.1 % triamcinolone acetonide in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxypropanoic acid 4.5 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. Triamcinolone acetonide 2% solution of acetone:ethanol (50:50), 5 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

EXAMPLE 45

10

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxypropanoic acid and 0.2 % 5-fluorouracil in a cream form for dermatologic disorders may be formulated as follows.

2-Hydroxypropanoic acid 4.5 g and creatinine 5.7 g are dissolved in 20 ml of water, and the solution thus obtained is mixed with 50 g of an oil-in-water emulsion. 5-Fluorouracil 2% solution of propylene glycol:water (95:5), 10 ml is added to the above mixture. More oil-in-water emulsion is added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

EXAMPLE 46

20

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 2% clotrimazole in a cream form for athlete's foot and other fungal infections may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g, clotrimazole 2 g and creatinine 5.7 g are dissolved in water 20 ml and propylene glycol 5 ml, and the solution thus obtained is mixed with enough amount of an oil-in-water emulsion to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

EXAMPLE 47

30

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 2% erythromycin in solution form for acne may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g, erythromycin 2 g and creatinine 5.7 g are dissolved in water 25 ml, ethanol 40 ml and propylene glycol 15 ml. More water is then added to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

EXAMPLE 48

40

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 1 % ketoconazole in a cream form for fungal infections may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g, ketoconazole 1 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with enough amount of an oil-in-water emulsion to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.2.

EXAMPLE 49

50

A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxypropanoic acid and 2% clotrimazole in a cream form for fungal infections may be formulated as follows.

2-Hydroxypropanoic acid 3.8 g, clotrimazole 2 g and creatinine 5.7 g are dissolved in 25 ml of water, and the solution thus obtained is mixed with enough amount of an oil-in-water emulsion to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated has pH 4.1.

EXAMPLE 50

5 A pseudoamphoteric composition containing in combination 0.5 M 2-hydroxyethanoic acid and 2% tetracycline in a gel form for dermatologic disorders may be formulated as follows.

2-Hydroxyethanoic acid 3.8 g, tetracycline 2 g, creatinine 5.7 g, xanthan gum 0.2 g, carbomer-941 1 g, propylene glycol 5 ml, ethanol 20 ml and enough amount of water are homogenized to make a total volume of 100 ml. The pseudoamphoteric composition thus formulated for acne and oily skin has pH 4.2.

10

EXAMPLE 51

15 An amphoteric composition containing 0.2 M aleuritic acid and 0.1 M L-lysine in a solution form for cosmetic and dermatologic conditions may be formulated as follows.

Aleuritic acid 6.1 g and L-lysine 1.5 g are dissolved in sufficient amount of a solution from ethanol:propylene glycol 80:20 to make a total volume of 100 ml. The amphoteric composition thus formulated has pH 6.4.

20

EXAMPLE 52

25 A typical composition containing a dimeric form of alpha hydroxyacid in solution for acne, dandruff, and as a skin cleanser may be formulated as follows.

Glycolide powder 1.0 g is dissolved in ethanol 89 ml and propylene glycol 10 ml. The composition thus formulated has pH 4.0, and contains 1% active ingredient.

30

EXAMPLE 53

A typical composition containing a dimeric form of alpha hydroxyacid in ointment for dry skin, psoriasis, eczema, pruritus, wrinkles and other skin changes associated with aging may be formulated as follows.

35 Glycolide powder 2.0 g is mixed uniformly with petrolatum 66 g and mineral oil 32 g. The composition thus formulated contains 2% active ingredient.

40

EXAMPLE 54

A typical composition containing a full strength or a high concentration of an alpha hydroxyacid, alpha ketoacid or closely related compound for topical treatments of warts, keratoses, acne, age spots, nail infections, wrinkles and aging related skin changes may be prepared as follows.

45 If the alpha hydroxyacid, alpha ketoacid or closely related compound at full strength is a liquid form at room temperature such as 2-hydroxypropanoic acid, 2-ketopropanoic acid, methyl 2-ketopropanoate and ethyl 2-ketopropanoate, the compound is directly dispensed as 0.5 to 1 ml aliquots in small vials. If the compound is a solid form at room temperature such as 2-hydroxyethanoic acid and 2-methyl 2-hydroxypropanoic acid, it is first dissolved in minimal amount of an appropriate solvent or solvent system such as water or ethanol and propylene glycol with or without a gelling agent. For example, 2-hydroxyethanoic acid 50 70 g is dissolved in water 30 ml, and the 70% strength 2-hydroxyethanoic acid thus obtained is dispensed as 0.5 to 1 ml aliquots in small vials. If a gelling agent is used, methyl cellulose or hydroxyethyl cellulose 1 g may be added to the above solution.

55

EXAMPLE 55

A typical composition containing an intermediate strength of an alpha hydroxyacid, alpha ketoacid or closely related compound for topical treatment of warts, keratoses, acne, nail infections, age spots, wrinkles and aging related skin changes may be prepared as follows.

2-Hydroxyethanoic acid or 2-ketopropanoic acid 40 g is dissolved in ethanol 54 g and propylene glycol 6 g, and the 40% strength solution thus obtained is dispensed as 5 to 10 ml aliquots in dropper bottles.

TEST RESULTS

10

In order to determine whether amphoteric and pseudoamphoteric compositions of the instant invention were therapeutically effective for various cosmetic conditions and dermatologic disorders, a total of more than 90 volunteers and patients participated in these studies. Some participating subjects were given two preparations; an amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound, and a vehicle placebo. Others were given multiple preparations containing a known pharmaceutical agent such as a corticosteroid with or without incorporation of an amphoteric or pseudoamphoteric composition consisting of an alpha hydroxyacid or the related compound of the instant invention. The amphoteric and pseudoamphoteric compositions were formulated according to the Examples described in the previous section.

20

1. Common dry skin.

Human subjects having ordinary dry skin or with moderate degrees of dry skin as evidenced by dryness, flaking and cracking of the skin were instructed to apply topically the lotion, cream or ointment containing an alpha hydroxyacid or the related compound in amphoteric or pseudoamphoteric composition, on the affected area of the skin. Topical application, two to three times daily, was continued for two to four weeks.

In all the 28 subjects tested, the feeling of the skin dryness disappeared within a week of topical application. The rough and cracked skin became less pronounced and the skin appeared normal and felt smooth after several days of topical treatment. The alpha hydroxyacids and the related compounds which have been found to be therapeutically effective when incorporated into the amphoteric or pseudoamphoteric compositions for dry skin are as follows:

2-hydroxyethanoic acid (glycolic acid), 2-hydroxypropanoic acid (lactic acid), 2-methyl-2-hydroxypropanoic acid (methylactic acid), phenyl 2-hydroxyethanoic acid (mandelic acid), phenyl 2-methyl-2-hydroxyethanoic acid (atrolactic acid), 3-phenyl-2-hydroxypropanoic acid (phenyllactic acid), diphenyl 2-hydroxyethanoic acid (benzilic acid), gluconolactone, tartaric acid, citric acid, saccharic acid, malic acid, tropic acid, glucuronic acid, galacturonic acid, gluconic acid, 3-hydroxybutanoic acid, quinic acid, ribonolactone, glucuronolactone, galactonolactone, pyruvic acid, methyl pyruvate, ethyl pyruvate, phenylpyruvic acid, benzoylformic acid and methyl benzoylformate.

The ordinary dry skin conditions, once restored to normal appearing skin, remained improved for some time until causes of dry skin, such as low humidity, cold weather, excessive contact pressure, detergents, soaps, solvents, chemicals, etc., again caused recurrence of the dry skin condition. On continued use it was also found that twice daily topical application of an amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound of the instant invention prevented the development of new dry skin lesions.

2. Severe dry skin.

50

In severe dry skin, the skin lesions are different from the ordinary dry skin. A main cause of severe dry skin is inherited genetic defects of the skin. The involved skin is hyperplastic, fissured and has thick adherent scales. The degree of thickening is such that lesions are palpably and visually elevated. The thickened adherent scales cause the surface of involved skin to be markedly rough and uneven. These two attributes of thickness and texture can be quantified to allow objective measurement of degree of improvement from topically applied test materials as follows:

55

DEGREE OF IMPROVEMENT					
	None	Mild	Moderate	Substantial	Complete
	(0)	(1 +)	(2 +)	(3 +)	(4 +)
Thickness	Highly elevated	Detectable reduction	Readily apparent reduction	Barely elevated	Normal thickness
Texture	Visibly rough	Palpably rough	Uneven but not rough	Slightly uneven	Visibly and palpably smooth

By means of such parameters, degrees of change in lesions can be numerically recorded and comparisons made of one treated site to another.

In order to evaluate the amphoteric and pseudoamphoteric compositions of the instant invention, a total of 6 patients having severe dry skin conditions were treated with the compositions containing an alpha hydroxyacid or the related compound.

Tested areas were of a size convenient for topical applications, i.e., circles 5 cm in diameter demarcated with a plastic ring of that size inked on a stamp pad. The medicinal lotions or creams were topically applied by the patient in an amount sufficient to cover the treatment sites. Applications were made three times daily and without occlusive dressings. Applications were discontinued at any time when resolutions of the lesion on the treatment area was clinically judged to be complete.

The test results of amphoteric and pseudoamphoteric compositions containing the following alpha hydroxyacids or the related compounds on patients with severe dry skin are summarized as follows:

4+ Effectiveness; glycolic acid, lactic acid, methylactic acid, mandelic acid, tropic acid, atrolactic acid and pyruvic acid.

3+ Effectiveness; benzoic acid, gluconolactone, malic acid, tartaric acid, citric acid, saccharic acid, methyl pyruvate, ethyl pyruvate, phenylactic acid, phenylpyruvic acid, glucuronic acid and 3-hydroxybutanoic acid.

2+ Effectiveness; mucic acid, ribonolactone, 2-hydroxydodecanoic acid, quinic acid, benzoylformic acid and methyl benzoylformate.

3. Psoriasis.

The involved skin in psoriasis is hyperplastic (thickened), erythematous (red or inflamed), and has thick adherent scales. The degree of thickening is such that lesions are elevated up to 1 mm above the surface of adjacent normal skin; erythema is usually an intense red; the thickened adherent scales cause the surface of involved skin to be markedly rough and uneven. These three attributes of thickness, color and texture can be quantified to allow objective measurement of degree of improvement from topically applied test materials as follows.

DEGREE OF IMPROVEMENT					
	None	Mild	Moderate	Substantial	Complete
	(0)	(1 +)	(2 +)	(3 +)	(4 +)
THICKNESS	Highly elevated	Detectable reduction	Readily apparent reduction	Barely elevated	Normal thickness
TEXTURE	Visibly rough	Palpably rough	Uneven but not rough	Slightly uneven	Visibly and palpably smooth
COLOR	Intense Red	Red	Dark Pink	Light Pink	Normal Skin Color

15	-----	-----
	Compositions*	Therapeutic Effectiveness
	-----	-----
20	Hydrocortisone 2.5% alone	1+
	With lactic acid	2+
	With glycolic acid	2+

	Compositions*	Therapeutic Effectiveness
5	With ethyl pyruvate	2+
	With methyl pyruvate	2+
10	With benzilic acid	2+
	With pyruvic acid	2+
	With methyllactic acid	2+
15	Hydrocortisone 17-valerate 0.2% alone	2+
	With lactic acid	3+
20	With glycolic acid	3+
	With benzilic acid	3+
	With ethyl pyruvate	3+
25	With methyl pyruvate	3+
	With gluconolactone	3+
30	With pyruvic acid	3+
	Betamethasone dipropionate 0.05% alone	3+
	With lactic acid	4+
35	With glycolic acid	4+
	With ethyl pyruvate	4+
40	With methyl pyruvate	4+
	With mandelic acid	4+
	With benzilic acid	4+
45	Clobetasol propionate 0.05% alone	3+
	With lactic acid	4+
50	With glycolic acid	4+

 Compositions*

 Therapeutic
Eff ctiveness

5	With ethyl pyruvate	4+
	With methyl pyruvate	4+
10	With methyllactic acid	4+
	With mandelic acid	4+
	With tropic acid	4+
15	With benzilic acid	4+

20 * Except the "alone" preparations, all others were
amphoteric or pseudoamphoteric compositions containing
0.2 to 2M alpha hydroxyacids or related compounds.

25

We have also found that an amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound in combination with an antimetabolite agent such as 5-fluorouracil with or without additional incorporation of a corticosteroid is therapeutically effective for topical treatment of
30 psoriasis.

4. Eczema.

35 In a topical treatment of eczema patients, hydrocortisone alone at 2.5% or hydrocortisone 17-valerate alone at 0.2% would achieve only 2+ improvement, and betamethasone dipropionate or clobetasol propionate alone at 0.05% would achieve only a 3+ improvement on all the eczema patients tested. Test results of amphoteric and pseudoamphoteric compositions containing both a corticosteroid and one of the following alpha hydroxyacids or the related compounds are shown as follows:

40 3+ Effectiveness; hydrocortisone 2.5% or hydrocortisone 17-valerate 0.2% plus lactic acid, glycolic acid, mandelic acid, ethyl pyruvate, gluconolactone, benzilic acid or ribonolactone.
4+ Effectiveness; betamethasone dipropionate or clobetasol propionate 0.05% plus lactic acid, glycolic acid, mandelic acid, ethyl pyruvate, methyl pyruvate, benzilic acid, gluconolactone, citric acid, tartaric acid or methyllactic acid.

45

5. Oily Skin and Skin Cleanse.

50 Human subjects having oily skin or blemished skin as well as acne patients having extremely oily skin participated in this study. Amphoteric and pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds were formulated in solution or gel form.

Each participating subject received a solution or a gel preparation containing an alpha hydroxyacid or a related compound in an amphoteric or pseudoamphoteric composition. The participating subjects were instructed to apply topically the solution or gel medication on the affected areas of forehead or other part of
55 the face. Three times daily applications were continued for 2 to 6 weeks.

The degree of improvement of oily skin as well as the rate of improvement of acne lesions were clinically evaluated. Most participants reported that oiliness of skin disappeared within one to two weeks of topical administration, and the skin so treated became smooth and soft. Many participating subjects

preferred gel preparations than solution compositions. It was found that all the participants showed substantial improvements on oily skin and acne lesions by six weeks of topical administration of amphoteric or pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds of the instant invention.

- 5 Those alpha hydroxyacids and the related compounds which have been found to be therapeutically effective for oily skin and as skin cleansers include: benzilic acid, glycolic acid, lactic acid, methylactic acid, mandelic acid, pyruvic acid, ethyl pyruvate, methyl pyruvate, tropic acid, malic acid, gluconolactone, 3-hydroxybutanoic acid, glycolide and polyglycolic acid. As a skin cleanser for oily skin or acne-prone skin, the amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound
10 may also be incorporated with other dermatologic agents. For example, an amphoteric gel composition may consist of both an alpha hydroxyacid and erythromycin or tetracycline.

6. Acne

- 15 Amphoteric and pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds of the instant invention in a solution or gel form were provided to patients having comedogenic and/or papulopustular lesions of acne. Each participating patient was instructed to apply topically the composition on the involved areas of the skin such as forehead, face and chest. Three times daily
20 administration was continued for 6 to 12 weeks.

- The degree and rate of improvement on acne lesions were clinically evaluated. It was found that acne lesions consisting mainly of comedones improved substantially after 6 to 8 weeks of topical administration with the amphoteric or the pseudoamphoteric composition containing an alpha hydroxyacid or the related compound. The time for complete clearing of comedogenic acne treated with the amphoteric or
25 pseudoamphoteric composition of the instant invention varied from 6 to 12 weeks.

- As a topical treatment for papulopustular and/or pustular acne the amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound may incorporate in addition an antiacne agent. The antiacne agents include antibiotics such as erythromycin, tetracycline, clindamycin, meclocycline and minocycline, and retinoids such as retinoic acid. Such combination compositions have
30 been found to be therapeutically more effective for topical treatment of severe acne.

7. Age Spots

- 35 Many small and large discolored lesions, commonly called age spots on the face and the back of the hands are benign keratoses, if they are not variants of actinic keratoses. Very few of such age spots are true lentigines, therefore alpha hydroxyacids and the related compounds may be effective in eradicating most age spots without concurrent use of skin bleaching agents such as hydroquinone and monobenzone. However, additional beneficial effects have been found when a skin bleaching agent such as hydroquinone
40 or monobenzone is also incorporated into the compositions of the instant invention for age spots involving pigmented lesions.

- Amphoteric and pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds, with or without incorporation of hydroquinone were provided to volunteer subjects and patients having age spot keratoses, melasma, lentigines and/or other pigmented lesions. Each participating subject
45 received two products, i.e., with or without the addition of 2% hydroquinone to the amphoteric or pseudoamphoteric composition containing an alpha hydroxyacid or the related compound.

- The volunteer subjects and patients were instructed to apply topically one medication on one side of the body such as left side of the face or on the back of the left hand, and the other medication on the other side of the body such as on right side of the face or on the back of the right hand. Specific instructions
50 were given to the participating subjects that the medications were applied three times daily to the lesions of age spot keratoses, melasmas, lentigines and/or other pigmented lesions. Clinical photos were taken of participating subjects before the initiation of the topical treatment and every 4 weeks during the course of treatment.

- At the end of 4 to 8 weeks, improvement of age spot keratoses was clinically discernible. After 4 to 6
55 months of topical treatment, substantial improvement of age spot keratoses occurred in the majority of subjects tested. Complete eradication of age spot keratoses occurred after 6 to 9 months of topical administration with the amphoteric or pseudoamphoteric compositions of the instant inventions.

Amphoteric or pseudoamphoteric compositions containing both an alpha hydroxyacid or the related

compound and hydroquinone were judged to be more effective in eradicating pigmented age spots, melasma, lentigines and other pigmented lesions.

The alpha hydroxyacids and the related compounds which have been found to be therapeutically effective for age spots with or without combination with hydroquinone include glycolic acid, lactic acid, methylactic acid, mandelic acid, pyruvic acid, methyl pyruvate, ethyl pyruvate, benzilic acid, gluconolactone, malic acid, tartaric acid, citric acid and tropic acid. For flat or slightly elevated seborrheic keratoses on the face and/or the back of the body, amphoteric or pseudoamphoteric compositions containing higher concentrations of alpha hydroxyacids or the related compounds have been found to be effective in eradicating such lesions.

Actinic keratoses may be successfully treated with amphoteric or pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds in combination with an antimetabolite agent such as 5-fluorouracil.

8. Warts.

Eradications of common warts by topical application of amphoteric or pseudoamphoteric compositions require higher than usual concentrations of alpha hydroxyacids or the related compounds in the formulations. The amphoteric or pseudoamphoteric compositions were formulated as a liquid or light gel form, and dispensed usually as 0.5-1 ml aliquots in small vials.

Topical applications were made discreetly to wart lesions by adult patients or by responsible adult family members. For ordinary usual warts of hands, fingers, palms and soles topical applications were made 2 to 4 times daily, and were continued for 2 to 6 weeks. Generally, the overlying stratum corneum of the wart lesion change in appearance after several weeks topical application of the composition. In most cases, the wart lesion simply fell off. The skin then healed normally without forming any scars.

We have also found that when a dermatologic agent such as 5-fluorouracil is incorporated into the amphoteric or pseudoamphoteric compositions containing alpha hydroxyacids or the related compounds, the medications have been very effective for topical treatment of warts without using higher concentrations of alpha hydroxyacids or the related compounds.

The alpha hydroxyacids and the related compounds which have been found to be therapeutically effective for topical treatment of warts with or without incorporation of 5-fluorouracil include glycolic acid, lactic acid, pyruvic acid, ethyl pyruvate, methyl pyruvate and mandelic acid.

Topical formulations and compositions containing specific alpha hydroxyacids, alpha ketoacids or the related compounds at full strengths or high to intermediate concentrations prepared according to Examples 54 and 55, without utilizing amphoteric or pseudoamphoteric systems, have also been tested for ordinary warts of the hands, fingers, palms and soles. Participating patients have been advised to apply a small drop of the medication with a toothpick or a fine caliber brush to the center of a wart lesion only. Prescribed applications have been 3 to 6 times daily, and are continued until the patient feels pain.

For the more rough-surfaced wart, the duration of application has been as short as one or a few days. For lesions with more compact, less permeable stratum corneum, the time to experience gain has been longer. Frequency and duration of applications have been modified according to other clinical responses and reactions of lesions, and the patient or responsible family member is instructed accordingly.

For example, some clinical manifestations other than pain have also been used as a signal to interrupt application. These manifestations have included distinct blanching of the lesions or distinct peripheral erythema. Very often, discomfort is the usual signal of clinical reactions.

Generally, the overlying stratum corneum of the wart lesions became loose, and the whole wart lesion simply fell off. The skin then healed normally without forming any scars.

9. Athlete's Foot and Nail Infections

Amphoteric and pseudoamphoteric compositions containing both an antifungal agent and one of the alpha hydroxyacids or the related compounds were provided to patients having frequent recurrence of fungal infections involving the foot. The antifungal agents include clotrimazole, miconazole, ketoconazole and griseofulvin. When both feet but not toe nails were involved in the infection, the patients were instructed to apply topically the compositions of the instant invention on the left foot, and a brand-name antifungal product on the right foot. Three times daily applications were continued for one to four weeks. The degree and rate of improvement on skin lesions were clinically evaluated, and comparison was made one side of

the body against the other. It was found that the skin lesions improved much faster with the amphoteric or pseudoamphoteric compositions containing both the antifungal agent and the alpha hydroxyacid or the related compound. The alpha hydroxyacids or the related compounds seemed to enhance the efficacies of the antifungal agents, and also to eliminate the discomforts such as itching, tingling, burning and irritation due to fungal infections. When toe nails were not involved the infected skin generally healed within one to two weeks from topical application of the amphoteric or pseudoamphoteric composition containing both an antifungal agent and an alpha hydroxyacid or the related compound.

Fungal infections of the nails are very difficult to treat, because antifungal products to date are not therapeutically effective for topical treatment of nails. One of the reasons is that most antifungal drugs have not been formulated as bioavailable forms in the commercial products. When tow nails were involved in the infections, patients were provided with amphoteric or pseudoamphoteric compositions containing in combination an antifungal agent and an alpha hydro xyacid or an alpha ketoacid at higher concentrations ranging from 20 to 99%, dispensed as 1-2 ml aliquots in small vials. The patients were instructed to apply topically the compositions discreetly to the infected nail surface by means of a fine calibre paint brush. the technique was the same as for application of nail polish, that is careful avoidance of contact with lateral nail folds or any peri-ungual skin. Once or twice daily applications were continued for 2 to 8 weeks.

As mentioned above, while brand-name antifungal products are usually not effective against fungus infections within or underneath the nail, it was found that the amphoteric or pseudoamphoteric compositions containing an antifungal agent and an alpha hydroxyacid or alpha ketoacid were therapeutically effective in eradicating fungal infections of the nails. Such treatment may cause in some instances the treated nail plate to become loose and eventually fell off from the nail bed. This happened quite naturally without any feeling of pain nor bleeding, and the skin lesion healed quickly with normal growth of a new nail.

10. Wrinkles

Wrinkles of skin may be due to natural aging and/or sun damage. Most fine wrinkles on the face are due to natural or innate aging, while coarse wrinkles on the face are the consequence of actinic or sun damage. Although the real mechanism of wrinkles formation in the skin is still unknown, it has been shown that visible fine wrinkles are due to diminution in the number and diameter of elastic fibers in the papillary dermis, and also due to atrophy of dermis as well as reduction in subcutaneous adipose tissue. Histopathology and electron microscopy studies indicate that coarse wrinkles are due to excessive deposition of abnormal elastic materials in the upper dermis and thickening of the skin. At present there are no commercial products which have been found to be therapeutically effective for topical eradication of wrinkles, although retinoic acid (tretinoin) has been shown to be beneficial for sun damaged skin.

In order to determine whether the amphoteric or pseudoamphoteric composition containing the alpha hydroxyacids, alpha ketoacids or the related compounds are therapeutically effective for wrinkles, patients and volunteer subjects participated in this study. The participants were instructed to apply the formulations of the instant invention twice daily on areas of facial wrinkles for 4 to 12 months. All participants were told to avoid sun exposure, and to use sunscreen products if exposure to sunlight was unavoidable. Photographs of each side of the face for each participant were taken at the beginning of the study and repeated at one to three-month intervals. The participants were asked not to wear any facial make-up at the time of each office visit. Standardized photographic conditions were used including the use of same lot of photographic film, the same light source at two feet from the face, aimed at a locus on the frontal aspect of each cheek. Each time photographs were taken with camera aimed perpendicular to the cheek. At the end of study twenty two participants had been entered into the study for at least four months. Clinical evaluations and review of photographs have revealed substantial reductions in facial wrinkles of the temporal region and cheek area on at least one side of the face in eighteen cases. Degree of improvement and reduction in wrinkles has been evaluated and determined to be mild to moderate in six participants but very substantial in twelve participants.

The alpha hydroxyacids, alpha ketoacids and other related compounds including their lactone forms which may be incorporated into the amphoteric and pseudoamphoteric compositions for cosmetic conditions and dermatologic disorders such as dry skin, acne, age spots, keratoses, warts and skin wrinkles or in combination with other dermatologic agents to enhance therapeutic effects include the following:

(1) Alkyl Alpha Hydroxyacids

2-Hydroxyethanoic acid (Glycolic acid), 2-Hydroxypropanoic acid (Lactic acid), 2-Methyl 2-hydroxypropanoic acid (Methyl lactic acid), 2-Hydroxybutanoic acid, 2-Hydroxypentanoic acid, 2-Hydroxyhexanoic acid, 2-Hydroxyheptanoic acid, 2-Hydroxyoctanoic acid, 2-Hydroxynonanoic acid, 2-Hydroxydecanoic acid, 2-Hydroxyundecanoic acid, 2-Hydroxydodecanoic acid (Alpha hydroxylauric acid), 2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid), 2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid), 2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid), 2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid).

(2) Aralkyl And Aryl Alpha Hydroxyacids

2-Phenyl 2-hydroxyethanoic acid (Mandelic acid), 2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid), 3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid), 2-Phenyl 2-methyl 2-hydroxyethanoic acid (Atrolactic acid), 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid, 2-(4'-Chlorophenyl) 2-hydroxyethanoic acid, 2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic acid, 2-(4'-Hydroxy-3'-methoxyphenyl) 2-hydroxyethanoic acid, 3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid, 3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid, 2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid.

(3) Polyhydroxy Alpha Hydroxyacids

2,3-Dihydroxypropanoic acid (Glyceric acid), 2,3,4-Trihydroxybutanoic acid (Isomers; erythronic acid, threonic acid), 2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic acid, arabinoic acid, xylonic acid, lyxonic acid), 2,3,4,5,6-Pentahydroxyhexanoic acid (Isomers; aldonic acid, altronic acid, gluconic acid, mannoic acid, gulonic acid, idonic acid, galactonic acid, talonic acid), 2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid, galactoheptonic acid, etc.)

(4) Polycarboxylic Alpha Hydroxyacids

2-Hydroxypropane 1,3-dioic acid (Tartronic acid), 2-Hydroxybutane-1,4-dioic acid (Malic acid), 2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid), 2-Hydroxy-2-carboxypentane-1,5-dioic acid (Citric acid), 2,3,4,5-Tetrahydroxyhexane-1,6-dioic acid (Isomers; saccharic acid, mucic acid, etc.)

(5) Alpha Hydroxyacid Related Compounds

Ascorbic acid, quinic acid, isocitric acid, tropic acid, 3-chlorolactic acid, trethocanic acid, cerebronic acid, citramalic acid, agaric acid, 2-hydroxynervonic acid and aleuritic acid.

(6) Alpha Ketoacids And Related Compounds

2-Ketoethanoic acid (Glyoxylic acid), Methyl 2-ketoethanoate, 2-Ketopropanoic acid (Pyruvic acid), Methyl 2-ketopropanoate (Methyl pyruvate), Ethyl 2-ketopropanoate (Ethyl pyruvate), Propyl 2-ketopropanoate (Propyl pyruvate), 2-Phenyl-2-ketoethanoic acid (Benzoylformic acid), Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate), Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate), 3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid), Methyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate), 2-Ketobutanoic acid, 2-Ketopentanoic acid, 2-Ketohexanoic acid, 2-Ketoheptanoic acid, 2-Ketooctanoic acid, 2-Ketododecanoic acid, Methyl 2-ketooctanoate

The amphoteric and pseudoamphoteric compounds which may be incorporated into the compositions of the instant invention for cosmetic and dermatologic conditions include amino acids, peptides, polypeptides, proteins and the like compounds such as creatinine and creatine.

The dimeric and polymeric forms of alpha hydroxyacids and the related compounds which may be incorporated into the compositions of the instant invention include acyclic esters and cyclic ester; for example, glycolyl glycollate, lactyl lactate, glycolide, lactide, polyglycolic acid and polylactic acid.

1. A pharmaceutical or cosmetic composition comprising in combination an amphoteric or pseudoamphoteric agent and an alpha hydroxyacid, an alpha ketoacid or a related compound in a pharmaceutically acceptable vehicle for topical application.
- 5 2. A composition comprising a cosmetic or pharmaceutical agent in an amphoteric or pseudoamphoteric system comprising an alpha hydroxyacid, an alpha ketoacid or a related compound in a pharmaceutically acceptable vehicle for topical treatment of cosmetic conditions or medical disorders.
3. A composition according to claim 2 wherein said cosmetic or pharmaceutical agent is selected from agents that improve or eradicate age spots, keratoses and wrinkles; analgesics; anaesthetics; antiacne
10 agents; antibacterials; antiyeast agents; antifungal agents; antiviral agents; antidandruff agents; antidermatitis agents; antipruritic agents; antiemetics; antimotionsickness agents; antiinflammatory agents; antihyperkeratolytic agents, antidry skin agents; antiperspirants; antipsoriatic agents; antiseborrheic agents; hair conditioners and hair treatment agents; antiaging and antiwrinkle agents; antiasthmatic agents and bronchodilators; sunscreen agents; antihistamine agents; skin lightening agents; depigmenting agents; vitamins;
15 corticosteroids; tanning agents; hormones; retinoids; topical cardiovascular agents or dermatologicals.
4. A composition according to claim 2 wherein said cosmetic or pharmaceutical agent is selected from clotrimazole, ketoconazole, miconazole, griseofulvin, hydroxyzine, diphenhydramine, pramoxine, lidocaine, procaine, mepivacaine, hydroquinone, monobenzone, erythromycin, tetracycline, clindamycin, meclocycline, minocycline, naproxen, ibuprofen, theophylline, cromolyn, albuterol, retinoic acid, 13-cis retinoic acid,
20 hydrocortisone, hydrocortisone 21-acetate, hydrocortisone 17-valerate, hydrocortisone 17-butyrate, betamethasone dipropionate, triamcinolone acetonide, fluicinsonide, clobetasol propionate, benzoyl peroxide, crotamiton, propranolol, promethazine, vitamin A palmitate or vitamin E acetate.
5. A composition according to any preceding claim which includes an amphoteric or pseudoamphoteric agent selected from amino acids, dipeptides, polypeptides, proteins, imidazoline derivatives, lecithin
25 derivatives, related agents or metallic oxides.
6. A composition according to any preceding claim which includes an amphoteric or pseudoamphoteric agent selected from glycine, alanine, valine, leucine, isoleucine, serine, threonine, cysteine, cystine, methionine, aspartic acid, asparagine, glutamic acid, glutamine, arginine, lysine, 5-hydroxylysine, histidine, phenylalanine, tyrosine, tryptophan, 3-hydroxyproline, 4-hydroxyproline, proline, homocysteine, homocystine,
30 homoserine, ornithine, citrulline, creatine, creatinine, 3-aminopropanoic acid, 2-aminobutanoic acid, 4-aminobutanoic acid, 2-amino-2-methylpropanoic acid, 2-methyl-3-aminopropanoic acid, theanine, phenylglycine, canavanine, cananine, 4-hydroxyarginine, 4-hydroxyornithine, homoarginine, 4-hydroxyhomoarginine, β -lysine, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid, 2,6-diaminopimelic acid, 2-amino-3-phenylbutanoic acid, 2-methylserine, 3-phenylserine, betaine, taurine, cysteinesulfinic acid, methionine sulfoxide,
35 methionine sulfone, 3,5-diiodotyrosine, thyroxine, monoiodotyrosine, pipecolic acid, 4-aminopipecolic acid, 4-methylproline, glycyglycine, carnosine, anserine, ophidine, homocarnosine, β -alanyllysine, β -alanylarginine, glutathione, ophthalmic acid, norophthalmic acid, bradykinin, glucagon, protamines, histones, cocoamphoglycine, cocoamphopropionate, cocamphopropylsulfonate, phosphatidyl ethanolamine, phosphatidyl serine, sphingomyelin, stearamidoethyl, diethylamine, stearamidoethyl diethanolamine,
40 stearamidopropyl dimethylamine, quaternary ammonium hydroxide, quaternium hydroxide, aluminum oxide or zinc oxide.
7. A composition according to any preceding claim wherein said alpha hydroxyacid is selected from alkyl alpha hydroxyacids, aralkyl and aryl alpha hydroxyacids, polyhydroxy alpha hydroxyacids and polycarboxylic alpha hydroxyacids having the following chemical formula:
45 (Ra) (Rb) C (OH) COOH
wherein Ra and Rb are H, F, Cl, Br, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and in addition Ra and Rb may carry OH, CHO, COOH and alkoxy group having 1 to 9 carbon atoms, said alpha hydroxyacid existing as a free acid or lactone form, or in salt form with an organic base or an inorganic alkali, and as
50 stereoisomers as D, L, and DL forms when Ra and Rb are not identical.
8. A composition according to claim 7 wherein said alkyl alpha hydroxyacid is selected from 2-Hydroxyethanoic acid (Glycolic acid), 2-Hydroxypropanoic acid (Lactic acid), 2-Methyl 2-hydroxypropanoic acid (Methylactic acid), 2-Hydroxybutanoic acid, 2-Hydroxypentanoic acid, 2-Hydroxyhexanoic acid, 2-Hydroxyheptanoic acid, 2-Hydroxyoctanoic acid, 2-Hydroxynonanoic acid, 2-Hydroxydecanoic acid, 2-Hydroxyundecanoic acid, 2-Hydroxydodecanoic acid (Alpha hydroxylauric acid), 2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid), 2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid), 2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid), 2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid).
55
9. A composition according to claim 7 wherein said aralkyl and aryl alpha hydroxyacid is selected from 2-

Phenyl 2-hydroxyethanoic acid (Mandelic acid), 2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid), 3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid), 2-Phenyl 2-methyl 2-hydroxyethanoic acid (Atrolactic acid), 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid, 2-(4'-Chlorophenyl) 2-hydroxyethanoic acid, 2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic acid, 2-(4'-Hydroxy-3'-methoxyphenyl) 2-hydroxyethanoic acid, 3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid, 3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid, or 2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid.

10. A composition according to claim 7 wherein said polyhydroxy alpha hydroxyacid and polycarboxylic alpha hydroxyacid is selected from 2,3-Dihydroxypropanoic acid (Glyceric acid), 2,3,4-Trihydroxybutanoic acid (Isomers; erythronic acid, threonic acid), 2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic acid, arabinonic acid, xylonic acid, lyxonic acid), 2,3,4,5,6-Pentahydroxyhexanoic acid (Isomers; allonic acid, altronic acid, gluconic acid, mannoic acid, gulonic acid, idonic acid galactonic acid, talonic acid), 2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid, galactoheptonic acid etc.), 2-Hydroxypropane-1,3-dioic acid (Tartronic acid), 2-Hydroxybutane-1,4-dioic acid (Malic acid), 2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid), 2-Hydroxy-2-carboxypentane-1,5-dioic acid (Citric acid), 2,3,4,5-Tetrahydroxyhexane-1,6-dioic acid (Isomers; saccharic acid, mucic acid, etc.), or lactone forms (gluconolactone, galactonolactone, glucuronolactone, galacturonolactone, gulonolactone, ribonolactone, saccharic acid lactone, pantoyllactone, glucoheptonolactone, mannnonolactone, galactoheptonolactone, etc.).

11. A composition according to any preceding claim wherein said alpha ketoacid has the following chemical formula:

20 (Ra) CO COO (Rb)

wherein Ra and Rb each represent H or an alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and in addition Ra may carry F, Cl, Br, I, OH, CHO, COOH or an alkoxy group having 1 to 9 carbon atoms, said alpha ketoacid existing as a free acid or an ester form, or in a salt form with an organic base or an inorganic alkali.

25 12. A composition according to claim 11 wherein said alpha ketoacid and its ester is selected from 2-Ketoethanoic acid (Glyoxylic acid), Methyl 2-ketoethanoate, 2-Ketopropanoic acid (Pyruvic acid), Methyl 2-ketopropanoate (Methyl pyruvate), Ethyl 2-ketopropanoate (Ethyl pyruvate), Propyl 2-ketopropanoate (Propyl pyruvate), 2-Phenyl-2-ketoethanoic acid (Benzoylformic acid), Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate), Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate), 3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid), Methyl 3-phenyl-2-ketopropanoate (Methyl phenylpyruvate), Ethyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate), 2-Ketobutanoic acid, 2-Ketopentanoic acid, 2-Ketohexanoic acid, 2-Ketopheptanoic acid, 2-Ketooctanoic acid, 2-Ketododecanoic acid, or Methyl 2-ketooctanoate.

30 13. A composition according to any preceding claim wherein the said related compound is selected from ascorbic acid, quinic acid, isocitric acid, tropic acid, trethocanic acid, 3-chlorolactic acid, cerebronic acid, citramalic acid, agaricic acid, 2-hydroxynervonic acid, aleuritic acid and pantoic acid.

14. A therapeutic composition for topical treatment of cosmetic conditions or dermatologic disorders comprising dimeric or polymeric forms of hydroxyacids, having the following chemical formula:

H [-O-C(Ra)(Rb)-CO-]_n OH

wherein Ra,Rb = H, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and n = 1 or any numerical number up to 200; Ra and Rb in monomer unit 2, 3, 4 may be the same or the different groups from that in monomer unit 1; the hydrogen atom in Ra and Rb may be substituted by a halogen atom or a radical of lower alkyl, aralkyl, aryl or alkoxy of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 9 carbon atoms, and the dimeric and polymeric forms of hydroxyacids may be present as a free acid, ester or in a salt form with an organic base or inorganic alkali.

45 15. A composition according to claim 14 wherein said dimeric or polymeric forms of hydroxyacids are selected from the group consisting of glycolyl glycollate, lactyl lactate, mandelyl mandellate, atrolactyl atrolactate, phenyllactyl phenyllactate, benzilyl benzillate, glycolyl lactate, lactyl glycollate, triglycolic acid, trilactic acid, polyglycolic acid or polylactic acid.

50 16. A therapeutic composition for topical treatment of cosmetic conditions or dermatologic disorders comprising dimeric or polymeric forms of hydroxyacids, having the following chemical formula:

[-O-C(Ra)(Rb)-CO-]_n

wherein Ra,Rb = H, alkyl, aralkyl or aryl group of saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms, and n = 1 or any numerical number, and Ra or Rb may be identical or not identical in the monomer units.

55 17. A composition according to claim 16 wherein said dimeric or polymeric forms of hydroxyacids are selected from glycolide, lactide, mandelide, atrolactide, phenyllactide, benzilide, methylactide, lactoglycolide or glycolactide.

18. A composition according to any one of claims 14 to 17 further comprising a cosmetic or pharmaceutical agent incorporated as an ingredient in said composition.

19. A composition according to any preceding claim for use in the treatment of dry skin, xerosis, ichthyosis, dandruff, brownish spots, keratoses, melasma, lentigines, age spots, liver spots, pigmented spots, wrinkles, blemishes, skin lines, oily skin, acne, warts, eczema, pruritic skin, psoriasis, inflammatory dermatoses, disturbed keratinization, skin changes associates with aging, nail or skin requiring cleansers, conditioning or treatment, and hair or scalp requiring shampooing or conditioning.

20. A therapeutic composition for topical treatment of warts, nail infections, age spots, wrinkles and aging related skin changes comprising at least one member selected from alpha hydroxyacids, alpha ketoacids or related compounds.

21. A composition according to claim 20 wherein said alpha hydroxyacids, alpha ketoacids or related compounds may be present as a free acid, lactone, ester or in salt form with an organic base or an inorganic alkali.

22. A composition according to claim 20 or claim 21 wherein said alpha hydroxyacids, alpha ketoacids or related compounds are selected from 2-hydroxyethanoic acid, 2-hydroxypropanoic acid, 2-methyl 2-hydroxypropanoic acid, 2-phenyl 2-hydroxyethanoic acid, 2,2-diphenyl 2-hydroxyethanoic acid, 2-phenyl 2-methyl 2-hydroxyethanoic acid, 2-phenyl 3-hydroxypropanoic acid, 2-ketopropanoic acid, methyl 2-ketopropanoate and ethyl 2-ketopropanoate.

23. A cosmetic skin treatment which comprises the topical application to the skin of a composition according to any preceding claim.

24. The use in the preparation of a pharmaceutical or cosmetic composition for the topical treatment of skin conditions, of a combination of an amphoteric or pseudoamphoteric agent and an alpha hydroxyacid, an alpha ketoacid or a related compound.

25

30

35

40

45

50

55



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 30 8828

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	EP-A-0 086 070 (UNILEVER NV) * Page 3, line 35 - page 4, line 20; page 7, lines 12-26 * - - -	1-24	A 61 K 7/48 A 61 K 31/19
X	LU-A-5 814 3 (MEDISAN AB) * Pages 8-9 in the whole (claims 1-8) * - - -	1-24	
A	EP-A-0 273 202 (E.J. VAN SCOTT) - - - - -	1-24	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			A 61 K
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of search 25 October 90	Examiner BRINKMANN C.
<div><div>CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</div><div>E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</div></div>			